

Table SI. Regional melanocytic naevi (MN) distribution at T1

Localization	Intervention n (%)	Control n (%)	Total n (%)
Missing data	12 (1.1)	12 (1.0)	24 (1.0)
Head excluding face	7 (0.6)	16 (1.3)	23 (1.0)
Face including ears	194 (17.2)	189 (15.1)	383 (16.1)
Neck, ventral	47 (4.2)	56 (4.5)	103 (4.3)
Neck, dorsal	48 (4.3)	62 (5.0)	110 (4.6)
Trunk, ventral	103 (9.1)	86 (6.9)	189 (8.0)
Trunk, dorsal	130 (11.5)	127 (10.2)	257 (10.8)
Upper arm, anterior	57 (5.1)	51 (4.1)	108 (4.5)
Upper arm, posterior	41 (3.6)	49 (3.9)	90 (3.8)
Lower arm, anterior	62 (5.5)	67 (5.4)	129 (5.4)
Lower with elbow, posterior	98 (8.7)	114 (9.1)	212 (8.9)
Thigh, anterior	96 (8.5)	124 (9.9)	220 (9.3)
Thigh, posterior	20 (1.8)	30 (2.4)	50 (2.1)
Lower leg with knee, anterior	83 (7.4)	102 (8.2)	185 (7.8)
Lower leg, posterior	36 (3.2)	46 (3.7)	82 (3.5)
Palms	8 (0.7)	16 (1.3)	24 (1.0)
Back of hands	74 (6.6)	72 (5.8)	146 (6.1)
Dorsum of the feet	9 (0.8)	26 (2.1)	35 (1.5)
Plantar	1 (0.1)	4 (0.3)	5 (0.2)
Total	1,114 (46.9)	1,237 (52.1)	2,375

$p = 0.116$  ( $\chi^2$  test).