

Table SI. Values of discrimination index (DI) derived from dermoscopic images of lesions taken at the first and final visit and the follow-up period

Case	Sex/race	Location of affected nail	Age at first visit, years	DI at the first (→and final) visit	Follow-up period, months
1	F/J	Right big toe	5	0.1293→0.1233	42.5
2	M/J	Left ring finger	12	0.1157→0.0706	69.6
3	F/J	Right thumb	8	0.0782→0.0566	81.5
4	F/J	Right thumb	10	0.2181→0.0868	46.5
5	M/J	Right big toe	9	0.1012→0.0779	69.7
6	F/J	Left thumb	5	0.0799→0.0936	103.4
7	F/J	Right thumb	5	0.0983→0.1380	41.6
8	F/J	Right thumb	2	0.1509→0.1572	97.7
9	F/J	Left big toe	5	0.0486→0.0622	26.7
10	M/J	Right thumb	18	0.1091→0.1425	33.6
11	F/J	Right thumb	18	0.0848→0.1266	30.6
12	F/J	Right index finger	17	0.0997	–
13	F/J	Left big toe	5	0.1235	–
14	M/C	Unknown finger	13	0.0700→0.0291	48.0
15	M/C	Unknown finger	3	0.0523	–

F: female, M: male, J: Japanese, C: Caucasian.