

Table SI. Demographics and clinical characteristics of the study population (n = 137*)

Age, years, mean \pm standard deviation	62.3 \pm 15.8
< 50 years, n (%)	30 (22.1)
50–64 years, n (%)	39 (28.7)
\geq 65 years, n (%)	67 (49.3)
Sex, n (%)	
Women	54 (40.0)
Men	81 (60.0)
Primary tumor location, n (%)	
Head/Neck	29 (21.8)
Trunk	55 (41.4)
Upper limbs	18 (13.5)
Lower limbs	31 (23.3)
Tumour type, n (%)	
Nodular malignant melanoma	39 (29.1)
Superficial spreading melanoma	63 (47.0)
Lentigo malignant melanoma	7 (5.2)
Other malignant melanoma	25 (18.7)
Breslow, mm, mean \pm standard deviation	2.3 \pm 2.7
Thin, n (%)	51 (39.5)
Intermediate, n (%)	59 (45.7)
Thick, n (%)	19 (14.7)
Ulceration, n (%)	
No	89 (74.8)
Yes	30 (25.2)
Metastatic malignant melanoma, n (%)	
No	86 (62.8)
Yes	51 (37.2)
Sentinel lymph node findings, n (%)	
Negative	56 (74.7)
Positive	19 (25.3)

*Numbers may not add-up to the total due to missing data.