

Another subentity of CSMH is characterized by a circumscribed annular patch with multiple follicular papules. Unlike classical CSMH, there is no hyperpigmentation and the hair pattern changes are less prominent (4).

There have also been reports of patients with multiple CSMH (5).

Diffuse CSMH has been described as part of the Michelin-tyre-baby syndrome (6). This syndrome is characterized by a combination of anomalies, such as ringed skin creases, cleft palate, epicanthal folds, hypertelorism, malformed ears and developmental delay (6).

On the basis of these criteria, we propose a clinical classification of CSMH of the skin.

1. Type 1: classical localized CSMH
2. Type 2: patchy follicular variant
3. Type 3: multiple CSMH
4. Type 4: diffuse CSMH

#### ACKNOWLEDGEMENT

We thank Mrs D. Steinbrecher and Mrs M. Bär for photography and Mrs K. Habenicht for her assistance with the literature search.

#### REFERENCES

1. Stokes JH. Nevus pilaris with hyperplasia of nonstriated muscle. *Arch Derm Syph* 1923; 7: 479–481.
2. Sourreil P, Beylot C, Delfour M. Hamartome par hyperplasie des muscles arrecteurs des poils chez un nourrisson d'un mois. *Bull Soc Fr Derm Syphilol* 1969; 76: 602.
3. Zvulunov A, Rotem A, Merlob P, Metzker A. Congenital smooth muscle hamartoma. *Am J Dis Child* 1989; 144: 782–784.
4. Tsambaos D, Orfanos CE. Cutaneous smooth muscle hamartoma. *J Cutan Pathol* 1982; 9: 33–42.
5. Guillot B, Huet P, Joujoux JM, Lorette G. Hamartomes musculaires lisses congénitaux multiples. *Ann Derm Venereol* 1999; 125: 118–120.
6. Schnur RE, Herzberg AJ, Spinner N, Kant JA, Magnusson M, McDonald-McGinn D, et al. Variability in the Michelin tyre baby syndrome. A child with multiple anomalies, smooth muscle hamartoma, and familial paracentric inversion of chromosome 7q. *J Am Acad Derm* 1993; 28: 364–370.

Accepted April 15, 1999.

Rainer Gerdson, Claudine Lagarde, Astrid Steen, Kay H. Steen, Manfred Uerlich and Thomas Bieber  
Department of Dermatology, University of Bonn, Sigmund Freud Strasse 25, D-53105 Bonn, Germany.

## Risk Factors for Skin Cancer in a Group of Renal Transplant Recipients

Sir,

Studies in northern Europe and Australia have detected a high incidence of skin cancer in renal transplant recipients (RTRs). Hartevelt et al. (1) found a cumulative incidence of skin cancer, ranging from 10% to 40%, at 10 and 20 years after transplantation. Bouwes Bavinck et al. (2), in a cohort of Australian patients, reported a higher incidence (from 45% to 70% at 10 and 20 years after transplantation), related to the more intense sun exposure at those latitudes. Only few reports have been published about RTRs from southern Europe and Italy. From 1990 to 1997 we enrolled a consecutive series of RTRs in a dermatological screening program, followed up by the Second Division of Surgery and Kidney Transplantation Centre of the Ospedale Civile Maggiore, Verona, Italy. All patients gave their informed consent before medical examination.

We examined 423 RTR subjects (290 males and 133 females), treated with 3 different immunosuppressive regimens: prednisolone+azathioprine (PA) (71 patients), prednisolone+azathioprine+cyclosporin (PAC) (191 patients), prednisolone+cyclosporin (PC) (161 patients). An accurate dermatological anamnesis was collected. Clinical records were available for all the patients. For each patient we recorded the following data: age, sex, stature, weight, date of the transplantation and of the visit, type and maintenance posology of the immunosuppressive drugs. The whole skin surface and mucous membranes were examined by a dermatologist. Any lesion suspected to be a skin cancer was excised and its histology examined. For estimates of potential prognostic factors, only variables available at the time of the transplant were considered: we took as an end point the diagnosis of cancer. Cox's proportional-hazards regression model (3),

performing backward stepping of variables with pre-assigned *p* values equal to 0.05 that controlled the stepping removal, and Kaplan and Meir survival curves were carried out.

The mean age of the RTRs at the visit was 46.2 years (SD 11.2, age range 19–68 years) and mean age at transplantation was 38.9 years (SD 11.7 age range 12–65 years), with no difference between males and females. Mean follow-up time was 7.6 years (SD 5.38, range 0–26 years): the PA group had a longer follow-up time (14.0 years) (SD 6.0) than the PAC group (6.6 years) (SD 3.9) and the PC group (5.9 years) (SD 4.7).

A total of 43 patients were excluded: 30 died of various accidents unrelated to the graft function and 13 rejected the graft (Table I).

A skin cancer was detected in 21/423 patients (5%) (BCC 10; Bowen 2; Keratoacanthomas 3; SCC 6). Cumulative incidence of the first cancer at 5, 10, 15 and 20 years of follow-up was 0.8%, 5.2%, 11.2% and 15.3%, respectively. The incidence of cancer was not significantly different in the 3

Table I. Patients excluded from the study

Immunosuppressive therapy	Death	Rejection	Total
PA	9	5	14
PAC	7	4	11
PC	14	4	18
Total	30	13	43

PA = prednisolone + azathioprine.

PAC = prednisolone + azathioprine + cyclosporin.

PC = prednisolone + cyclosporin.

