

xeroderma pigmentosum complementation group F and a morphoic basal cell carcinoma. *Br J Dermatol* 1995; 132: 122–127.

- Lambert WC, Lambert MW. Diseases associated with DNA and chromosomal instability. In Alper JC, editor. *Genetic diseases of the skin*. St. Louis: Mosby Year Book 1991: 320–358.
- Lambert WC, Kuo HR, Lambert MW. Xeroderma pigmentosum. *Derm Clin* 1995; (13) 1: 169–209.

Accepted July 30, 1998.

G. Tessari<sup>1</sup>, F. S. D'Onghia<sup>1</sup>, M. Stefanini<sup>2</sup> and A. Barba<sup>1</sup>  
<sup>1</sup>Department of Dermatology, University of Verona, c/o Ospedale Civile Maggiore Piazzale Stefani 1 37124 Verona and <sup>2</sup>Consiglio Nazionale delle Ricerche – Istituto di Genetica Biochimica ed Evoluzionistica, Pavia, Italy.

## Post-herpes Zoster Scar Sarcoidosis

Sir,

### CASE REPORT

A 70-year-old white woman presented a papular zosteriform eruption on her right upper thorax and right arm. Shiny, erythematous-purplish or flesh-coloured papules 2–3 mm in diameter were observed, isolated or closely arranged in little plaques (Fig. 1). The patient complained only of mild itching; no systemic signs and symptoms were referred. Two months earlier the patient had a C5–C6 right brachial palsy due to brachial herpes zoster. The histopathological pattern of a papula showed a dermal non-caseating granulomatous infiltrate with giant Langerhans-like cells and sclerosing evolution indicating scar sarcoidosis. Further clinical and laboratory investigations failed to reveal systemic sarcoidosis. Chest roentgenogram and liver ultrasonography were normal; X-ray of the hands and feet did not reveal osteolytic alterations. Spirometry and ophthalmology examinations were negative, as well as Mantoux test and skin tests.

### DISCUSSION

The infiltration of scar tissue by non-caseating granulomas is a well-recognized form of cutaneous sarcoidosis (1–3). To our knowledge, it has been reported to occur in the site of previous herpes zoster only in one case (1).

Most of the patients with scar sarcoidosis have other systemic manifestations, particularly pulmonary changes; scar infiltrates may appear early in the disease or before parenchymal changes. Changes in scars in patients with sarcoidosis in remission may indicate exacerbation of the disease or may even be a marker for recurrences of thoracic sarcoidosis (4). A careful and prolonged follow-up in these patients is strongly recommended due to the potential risk of developing systemic sarcoidosis.

### REFERENCES

- Bisaccia E, Scarborough DA, Carr RD. Cutaneous sarcoid granuloma formation in herpes zoster scars. *Arch Dermatol* 1983; 119: 788–789.



Fig. 1. Papular zosteriform eruption of the right arm.

- Olumide YM, Bandle EO, Elesha SO. Cutaneous sarcoidosis in Nigeria. *J Am Acad Dermatol* 1989; 21: 1222–1224.
- Veien NK, Stahl D, Brodthagen H. Cutaneous sarcoidosis in Caucasians. *J Am Acad Dermatol* 1987; 16: 534–540.
- Roegel E, Pauli G, Grosshans E, El Baz D, Dietemann-Molard A, Hutt JP. Sarcoides récidivantes sur cicatrices cutanées révélatrices de sarcoidose thoracique itérative. *Rev Pneumol Clin* 1984; 40: 81–83.

Accepted July 30, 1998.

Monica Corazza, Sandra Bacilieri and Renata Strumia  
 Dermatology Department, University of Ferrara, Via Savonarola 9,  
 I 44100 Ferrara, Italy.