

LETTER TO THE EDITOR

**Comment on Zhao et al. “Palmoplantar Keratoderma of the Gamborg-Nielsen Type is Caused by Mutations in the *SLURP1* Gene and Represents a Variant of Mal de Meleda”**

**Ruud G. L. Nellen<sup>1,3</sup>, Peter M. Steijlen<sup>2,3</sup>, Michel van Geel<sup>2,4</sup> and Maurice A. M. van Steensel<sup>5</sup>**

<sup>1</sup>Department of Dermatology, IJsselland Ziekenhuis, Prins Constantijnweg 2, NL-2906 ZC Capelle aan den IJssel, <sup>2</sup>Department of Dermatology, <sup>3</sup>GROW School for Oncology and Developmental Biology, and <sup>4</sup>Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, The Netherlands, and <sup>5</sup>Institute of Medical Biology, Immunus, Singapore. E-mail: rnellen@ysl.nl

Sir,

As reported recently in *Acta Dermato-Venereologica*, Zhao et al. (1) identified mutations in the *SLURP1* gene, encoding the secreted lymphocyte antigen 6/urokinase-type plasminogen activator receptor related protein-1 in patients with palmoplantar keratoderma of the Gamborg-Nielsen type (PPK-GN). They describe 15 patients with a marked, sharply demarcated, and waxy, usually yellowish PPK with an erythematous base, involvement of the dorsal sides of the fingers and toes, interspersions of erythematous skin in the hyperkeratosis and pseudo-ainhum. The clinical phenotype was complicated by hyperhidrosis and foul odour due to fungal superinfection. Symptoms began in the first year of life. The authors did not report skin changes on other parts of the body or mucosal symptoms. They suggest that, given the relatively mild phenotype, PPK-GN should be considered as a distinct entity allelic to Mal de Meleda (MDM) which is also caused by mutations in *SLURP1* (2).

We disagree with this suggestion. To classify a disease as allelic to another genetic disease caused by mutations in the same gene, the symptoms need to be distinct and (partly) non-overlapping, not demonstrate varying severity. For instance, both Papillon-Lefèvre syndrome (PLS, MIM #245000) and Haim-Munk syndrome (HMS, MIM #245010) are caused by mutations in the gene coding for cathepsin C (*CTSC*) (3). Their phenotypes overlap to include PPK and periodontitis, but in Haim-Munk syndrome, arachnodactyly, acroosteolysis, and onychogryphosis also develop. Because of this, HMS is considered a distinct entity allelic to PLS. Clearly, the situation for PPK-GN and MDM is quite different. These 2 conditions represent a spectrum of disease severity, as is seen in most genetic disorders. The cases described by Zhao et al. (1) are not remarkable for being especially mild or otherwise distinct from classic MDM.

To illustrate the point, we recently reported a Dutch cohort of MDM-patients with a phenotype comparable to the patients reported by Zhao et al. (1) All subjects had PPK with transgradient hyperkeratosis, pseudo-ainhum and hyperhidrosis with bacterial superinfection, with no skin symptoms elsewhere or mucosal changes (4). All patients had the p.Trp15Arg mutation in *SLURP1*, as was found in the majority of patients in the study by Zhao et al. (1). A relatively mild phenotype has been reported in a significant number of other studies on MDM, with varying severity of symptoms reported even within families harbouring the same mutation in *SLURP1* (5–9). Table S1<sup>1</sup> semi-quantitatively summarizes the phenotypes reported so far in individual patients with MDM, with a description of disease course, including age, clinical characteristics and *SLURP1* mutations. We added the data from the Dutch and Swedish cohorts. We are aware that drawing firm conclusions from these data is difficult; there are no reports describing large cohorts of patients and we acknowledge that there are no objective measures of MDM severity. That said, there seems to be a wide range of symptom severity, with previously reported cases of MDM being similar to the patients reported by Zhao et al. (1) The data suggest a tendency to a more severe course in older patients, which is consistent with earlier reports that note gradual progression during life. It would be of interest to follow disease progression and variation in a larger cohort of patients in order to obtain a more complete view of phenotypic variation in MDM.

In conclusion, we consider that PPK-GN represents only the mild end of the MDM spectrum. We therefore propose that the classification of PPK-GN as a separate genodermatosis should be abandoned.

<sup>1</sup><https://doi.org/10.2340/00015555-2196>

**Reply to Nellen et al’s Comment on the Classification of Clinical/genetic Variants of Mal de Meleda**

**Linshu Zhao<sup>1</sup>, Anders Vahlquist<sup>2\*</sup>, Marie Virtanen<sup>2</sup>, Lena Wennerstrand<sup>3</sup>, Lisbet K. Lind<sup>3</sup>, Anita Lundström<sup>4</sup> and Maritta Hellström Pigg<sup>1</sup>**

Departments of <sup>1</sup>Immunology, Genetics and Pathology and <sup>2</sup>Medical Sciences, Uppsala University, Uppsala, and <sup>3</sup>Departments of Medical Biosciences, Medical and Clinical Genetics and <sup>4</sup>Public Health and Clinical Medicine, Umeå University, Umeå, Sweden

We thank Nellen et al. (1) for their valuable comments on our recent paper in *Acta Dermato-Venereologica* in

which we showed for the first time that palmo-plantar keratoderma of the Gamborg-Nielsen type (PPK-GN, OMIM

244850) is caused by recessive mutations in the same gene as Mal de Meleda (MdM; OMIM 248300). Several pedigrees and clinical as well as ultrastructural characteristics of PPK-GN were described by Gamborg-Nielsen and co-workers in the late 1980s as a distinct disorder separate from autosomal dominant PPK, type Bothnia, which is very frequent in northern Sweden (18, 19). Until our recent publication, PPK-GN was considered by many as a distinct disease with as yet unknown aetiology. Our demonstration of recurrent and novel *SLURP-1* mutations in many of the Swedish patients with PPK-GN clearly identified this entity as a mild variant of MdM, indeed similar to what has been found in isolated cases of MdM

in several countries, including Holland (4, 6; see Table I).

In their commentary, Nellen et al. dispute our mentioning on the last line of the Discussion (1) that “PPK-GN and MdM indeed are allelic disorders”. We agree that this was pushing the distinction too far; perhaps a better phrasing would have been “PPK-GN is an allelic variant of MdM”. But PPK-GN is still described in many textbooks and in McKusick’s database (MIM) as a distinct entity; so, for both practical and historic reasons it may be of some value to use this eponym in parallel with MdM, at least in Sweden, where over the years many patients have been given the PPK-GN diagnosis based solely on clinical findings and inheritance pattern.

#### REFERENCES (for both papers)

- Zhao L, Vahlquist A, Virtanen M, Wennerstrand L, Lind L, Lundstrom A, et al. Palmoplantar keratoderma of the Gamborg-Nielsen type is caused by mutations in the *SLURP1* gene and represents a variant of Mal de Meleda. *Acta Derm Venereol* 2014; 94: 707–710.
- Fischer J, Bouadjar B, Heilig R, Huber M, Lefevre C, Jobard F, et al. Mutations in the gene encoding *SLURP-1* in Mal de Meleda. *Hum Mol Genet* 2001; 10: 875–880.
- Hart TC, Hart PS, Michalec MD, Zhang Y, Firatli E, Van Dyke TE, et al. Haim-Munk syndrome and Papillon-Lefevre syndrome are allelic mutations in cathepsin C. *J Med Genet* 2000; 37: 88–94.
- Nellen RG, Steijlen PM, Hennies HC, Fischer J, Munro CS, Jonkman MF, et al. Haplotype analysis in western European patients with mal de Meleda: founder effect for the W15R mutation in the *SLURP1* gene. *Br J Dermatol* 2013; 168: 1372–1374.
- Bchetnia M, Bozgia M, Laroussi N, Ben Brick AS, Charfeddine C, Ben Halim N, et al. The first Mal de Meleda case in Libya: identification of a *SLURP1* mutation. *Int J Dermatol* 2014 Apr 16. [Epub ahead of print].
- Eckl KM, Stevens HP, Lestringant GG, Westenberger-Treumann M, Traupe H, Hinz B, et al. Mal de Meleda (MDM) caused by mutations in the gene for *SLURP-1* in patients from Germany, Turkey, Palestine, and the United Arab Emirates. *Hum Genet* 2003; 112: 50–56.
- Zhang J, Cheng R, Ni C, Liang J, Li M, Yao Z. First Mal de Meleda report in Chinese Mainland: two families with a recurrent homozygous missense mutation in *SLURP-1*. *J Eur Acad Derm Venereol* 2015 Mar 12. [Epub ahead of print].
- Yerebakan O, Hu G, Yilmaz E, Celebi JT. A novel mutation in the *ARS* (component B) gene encoding *SLURP-1* in a family with Mal de Meleda. *Clin Exp Dermatol* 2003; 28: 542–544.
- Bchetnia M, Merdassi A, Charfeddine C, Mgaïeth F, Kassar S, Ouechtati F, et al. Coexistence of mal de Meleda and congenital cataract in a consanguineous Tunisian family: two case reports. *J Med Case Reports* 2010; 4: 108.
- Nellen RG, van Geel M, Steijlen PM, van Steensel MA. Compound heterozygosity for *ARS* component B mutations in a Dutch patient with mal de Meleda. *Br J Dermatol* 2009; 160: 878–880.
- Sakabe J, Kabashima-Kubo R, Kubo A, Sasaki T, Tokura Y. A Japanese case of Mal de Meleda with *SLURP1* mutation. *J Dermatol* 2014; 41: 764–765.
- Bchetnia M, Laroussi N, Youssef M, Charfeddine C, Ben Brick AS, Boubaker MS, et al. Particular Mal de Meleda phenotypes in Tunisia and mutations founder effect in the Mediterranean region. *BioMed Res Int* 2013; 2013: 206803.
- Musulmanoglu MH, Saracoglu N, Cilingir O, Basmaci T, Urer S, Sabuncu I, et al. A novel mutation in the *ARS* (component B) gene encoding *SLURP-1* in a Turkish family with mal de Meleda. *Br J Dermatol* 2006; 155: 467–469.
- Gruber R, Hennies HC, Romani N, Schmuth M. A novel homozygous missense mutation in *SLURP1* causing Mal de Meleda with an atypical phenotype. *Arch Dermatol* 2011; 147: 748–750.
- Chao SC, Huang CY, Lai FJ, Yang MH. Pseudodominant inheritance with the G86R mutation in the *ARS* gene in Mal de Meleda. *Int J Dermatol* 2006; 45: 1456–1458.
- Oh YJ, Lee HE, Ko JY, Ro YS, Yu HJ. A sporadic case of Mal de Meleda caused by gene mutation in *SLURP-1* in Korea. *Ann Dermatol* 2011; 23: 396–399.
- Taylor JA, Bondavalli D, Monif M, Yap LM, Winship I. Mal de Meleda in Indonesia: Mutations in the *SLURP1* gene appear to be ubiquitous. *Australas J Dermatol* 2014 Dec 30. [Epub ahead of print].
- Gamborg Nielsen P. Two different clinical and genetic forms of hereditary palmoplantar keratoderma in the northernmost county of Sweden. *Clin Genet* 1985; 28: 361–366.
- Kastl I, Anton-Lamprecht I, Gamborg Nielsen P. Hereditary palmoplantar keratosis of the Gamborg-Nielsen type. Clinical and ultrastructural characteristics of a new type of autosomal recessive palmolantar keratosis. *Arch Dermatol Res* 1990; 282: 363–370.