

DEBATE

HOW SHOULD WE USE THE VISUAL ANALOGUE SCALE (VAS) IN REHABILITATION OUTCOMES? IV: REPLY ON “HOW SHOULD WE USE THE VISUAL ANALOGUE SCALE (VAS) IN REHABILITATION OUTCOMES?”

Paula Kersten, PhD¹, Ayse A. Küçükdeveci, MD² and Alan Tennant, PhD³

From the ¹Person Centred Research Centre, School of Rehabilitation and Occupation Studies, Auckland University of Technology, Auckland, New Zealand, ²Department of Physical Medicine & Rehabilitation, Faculty of Medicine, Ankara University, Turkey and ³Department of Rehabilitation Medicine, Faculty of Medicine and Health, University of Leeds, UK. E-mail: pkersten@ait-ac-nz

We would like to respond to the commentaries given (1–3) to our review on Visual Analogue Scales (VAS), which concluded that there is sufficient evidence that VAS data are ordinal and that the VAS should be analysed appropriately from a statistical perspective by using non-parametric statistics (4). The three commentaries raise a number of important points and somewhat different perspectives. Franchignoni et al. (1) comment that, from a practical point of view, the VAS is harder to understand and that patients take longer to score it than a Likert scale (5); that the VAS has lower compliance rates (6); that it takes longer for the clinician to obtain the data (measuring by the clinician) and that there are possibilities of clinician measurement error. This concurs with previous findings that the VAS is difficult for some people (7) and has low test-retest reliability as a consequence (8). By contrast, Price et al. (2) contend that the VAS is easy to use, although they also agree that for some people it is not. They further say that the literature suggests the VAS has better psychometric properties than the Numeric Rating Scale (NRS). Harms-Ringdahl makes the point about the variability of the anchors, which may be used, and raises important issues about sample size, distribution and the relationship between pain and physical functioning (3).

We think it important to differentiate the question of the similarity (or lack) of the NRS and VAS, and various operational matters with respect to the ease of use and other practical matters associated with the VAS, from the primary question we tried to address, which was that of the scaling property of the VAS. In this respect, Price et al. (2) argue that the VAS has ratio properties because ‘separate judgments of ratios or proportions of pain intensity are in quantitative agreement with VAS ratings of pain’ and that studies ‘directed toward *both* experimental and clinical pain shows reliable stimulus-response power functions and a zero point for the VAS scale’ (9–13). In addition, they claim that if VAS has ratio properties it must also have interval properties. This is much the same point made by Harms-Ringdahl who refers to findings from her experimental pain study in which pain was measured on the VAS and the Borg Category Ratio scale (14). She states that in this study ‘pain intensity levels on both scales followed each other nicely’ and that ‘the non linear but logarithmic increase per se in pain intensity’ seems to be an invalid reason for our recommendation not to use VAS assessments as interval data. She further indicates that we have failed to describe experimental

studies where VAS has been found to allow measurements not only with interval but with ratio properties (3).

We would raise a number of points with respect to these assertions. Primarily, we would argue that the samples used in these studies, and the methods adopted do not allow for an assessment of ratio or, indeed, interval properties. In the original studies by Price et al. pain patients were subjected to noxious thermal stimuli to the forearm (11–13). As the power functions between VAS scores and the temperatures were predictive of estimated ratios of sensation or affect produced by pairs of standard temperatures (e.g. 47°C and 49°C), they concluded that this was evidence for ratio scaling properties of VAS.

We would suggest there are a number of issues with respect to this interpretation. Degree centigrade has only an arbitrary zero and is not a ratio scale in the first place. Only degree Kelvin can be said to be so. Thus deriving a ratio scale property using the measurement of degree centigrade cannot be undertaken (unless presumably the anchor for the VAS was –273 °C or their ‘reliable’ zero point is consistent with zero degree Kelvin). However, this does not preclude concluding that the scale is at the interval level, which itself is a necessary but not sufficient condition for ratio scaling.

Our primary concern relates to the fallacy of extrapolation from a narrow range, as typically undertaken in the studies reported by both Price et al. (2) and Harms-Ringdahl (3). The range of temperature extrapolation, from a few degrees to the whole, as in the Price et al. studies (11–13), represents a classical error. In part, this danger is recognised by the later findings of Myles et al. (9, 10). These studies asked patients with pain to score their pain and once it was halved (by means of pain relief) to score the VAS again and when they considered pain relief was satisfactory, the VAS was scored a third time (9). Similarly, patients were asked to consider what their pain might be like if it were twice as bad and score this on a VAS (10). The important point from this study is that the authors remind the reader that the sample had only mild-to-moderate pain and they explicitly state that the VAS may be non-linear at the margins. Thus they recognise the potential error of extrapolation from a narrow band in the centre.

In fact, the findings from these studies are quite consistent with information derived from Rasch analysis of the VAS, given the extremely limited range of stimuli presented to their subjects.

We would remind readers that the Rasch model is consistent with a probabilistic form of the theory of additive conjoint measurement, which is how data from ordinal scales can be tested for quantitative structure and interval scaling (15, 16). In their study, Price et al. (11) appear to have used a VAS of 150 mm in length, although they report that 100 is equal to the most intense sensation imaginable. Given a presumed rescaling to 0–100, they report the mean intensity values of pain at the ‘minimum’, ‘usual’ and ‘maximal’ levels during the previous week were 20.0, 49.3 and 72.3 °C, and 3 matching noxious temperatures as direct matches to their chronic pain at 43.1, 46.2 and 48.8 °C, respectively. Given their reported VAS scores were largely operating over the middle region of the scale, the empirical Rasch-based evidence does suggest that VAS will operate linearly (17). Outside of the middle range we have shown the VAS to be increasingly non-linear, such that points at the margins represent several magnitudes greater than a point in the middle region of the trait being measured (17). This was in fact the second such test of VAS data against the Rasch model, following an early analysis by Thomee et al. (18). In addition, traditional psychometric studies have also demonstrated that the VAS does not behave linearly (19–21).

Thus, in conclusion we would contend that the findings of the early experimental studies do not support interval scaling for the VAS and that, from first principles, given the experimental validation was based upon extrapolation from a narrow range of temperature, it cannot support the supposition of either an interval or ratio scale. Consequently we disagree with Price et al. (2) that there is insufficient evidence that the VAS is ordinal. They state strongly that our assertions of ordinality contain faulty assertions that could have destructive consequences for pain measurement. Rather, we would contend that holding on to the belief that the VAS is interval risks misinference, and could lead to flawed and inappropriate conclusions of trials, particularly where the levels of pain are outside the central portion of the VAS scale. Consequently, based on our review and research discussed above we maintain that parametric statistics should not be used for VAS rating data because the VAS only has ordinal scale properties. If Price and others wish to reject this evidence, they should refute Luce and Tukey’s theory, and the mathematical proofs that the Rasch model is consistent with that theory in a probabilistic manner (16, 22). Otherwise the empirical evidence is unequivocal; VAS is ordinal.

REFERENCES

1. Franchignoni F, Salaffi F, Tesio L. How should we use the visual analogue scale (VAS) in rehabilitation outcomes? I: How much of what? The seductive VAS numbers are not true measures. *J Rehabil Med* 2012; 44: 798–799.
2. Price DD, Staud R, Robinson ME. How should we use the visual analogue scale (VAS) in rehabilitation outcomes? II: Visual analogue scales as ratio scales: An alternative to the view of Kersten et al. *J Rehabil Med* 2012; 44: 800–801.
3. Harms-Ringdahl K. How should we use the visual analogue scale (VAS) in rehabilitation outcomes? III: On the validation requirements for assessments using VAS with ratio properties. *J Rehabil Med* 2012; 44: 801–802.
4. Kersten P, Kükükdveci AA, Tennant A. The use of the Visual Analogue Scale (VAS) in Rehabilitation Outcomes. *J Rehabil Med* 2012; 44: 609–610.
5. Guyatt G, Townsend M, Berman L, Keller J. A comparison of Likert and visual analogue scales for measuring change in function. *J Chronic Dis* 1987; 40: 1129–1133.
6. Hjermstad MJ, Fayers PM, Haugen DF, Caraceni A, Hanks GW, Loge JH, et al. Studies comparing numerical rating scales, verbal rating scales, and visual analogue scales for assessment of pain intensity in adults: a systematic literature review. *J Pain Symp Man* 2011; 41: 1073–1093.
7. Jackson D, Horn S, Kersten P, Turner-Stokes L. Development of a pictorial scale of pain intensity for patients with communication impairments: Initial validation in a general population. *Clin Med* 2006; 6: 580–585.
8. Carlsson AM. Assessment of chronic pain. I. Aspects of the reliability and validity of the visual analogue scale. *Pain* 1983; 16: 87–101.
9. Myles PS, Urquhart N. The linearity of the visual analogue scale in patients with severe acute pain. *Anaesthesia Intensive Care* 2005; 33: 54–58.
10. Myles PS, Troedel S, Boquest M, Reeves M. The pain visual analog scale: Is it linear or nonlinear? *Anesthesia Analgesia* 1999; 89: 1517–1520.
11. Price D, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain* 1983; 17: 45–56.
12. Price DD, Bush FM, Long S, Harkins SW. A comparison of pain measurement characteristics of mechanical visual analogue and simple numerical rating scales. *Pain* 1994; 56: 217–226.
13. Price DD, Harkins SW. Combined use of experimental pain and visual analogue scales in providing standardized measurement of clinical pain. *Clin J Pain* 1987; 3: 1–8.
14. Harms-Ringdahl K, Carlsson AM, Ekholm J, Raustorp A, Svensson T, Toresson HG. Pain assessment with different intensity scales in response to loading of joint structures. *Pain* 1986; 27: 401–411.
15. Rasch G. Probabilistic models for some intelligence and attainment tests (revised and expanded ed.). Chicago: The University of Chicago Press; 1960/1980.
16. Luce RD, Tukey JW. Simultaneous conjoint measurement: A new type of fundamental measurement. *J Math Psychol* 1964; 1: 1–27.
17. Kersten P, White PJ, Tennant A. The Visual Analogue WOMAC 3.0 scale – Internal validity and responsiveness of the VAS version. *BMC Musculoskeletal Dis* 2010; 11: 80.
18. Thomee R, Grimby G, Wright BD, Linacre JM. Rasch analysis of Visual Analog Scale measurements before and after treatment of Patellofemoral Pain Syndrome in women. *Scand J Rehabil Med* 1995; 27: 145–151.
19. Forrest M, Andersen B. Ordinal scale and statistics in medical research. *BMJ* 1986; 292: 537–538.
20. Munshi J. A method for constructing Likert scales [Internet]. Sonoma State University 1990 [cited 2011 Nov 7]. Available from: <http://www.munshi.4t.com/papers/likert.html>.
21. Svensson E. Guidelines to statistical evaluation of data from rating scales and questionnaires. *J Rehabil Med* 2001; 33: 47–48.
22. Van Newby A, Conner GR, Bunderson CV. The Rasch model and additive conjoint measurement. *J Appl Meas* 2009; 10: 348–354.

Submitted June 28, 2012; accepted June 29, 2012