

ERRATUM

In: von Bülow C, Amris K, la Cour K, Danneskiold-Samsøe B, Ejlersen Wæhrens E. Differences in ability to perform activities of daily living among women with fibromyalgia: A cross-sectional study. J Rehabil Med 2015; 47: 941–947.

Tables I–II are incorrect due to minor miscalculations; the conclusions are the same and the correct Tables are shown below.

Table I. Baseline characteristics in the overall study population and study participants subgrouped based on Assessment of Motor and Process Skills (AMPS) independence cut-offs

	Total	Group A	Group B	Group C	Group D	Group difference*	Significant differences between groups**		
						F; df; <i>p</i> -value	Group	Difference	<i>p</i> -value
Participants, <i>n</i> (%)	257 (100)	93 (36)	116 (45)	14 (6)	34 (13)				
Age, mean (SD)	45.39 (9.83)	46.11 (9.95)	46.18 (9.38)	47.65 (10.94)	39.76 (9.01)	422.72; 3; 0.004	A/B	0.07	1.00
							A/C	1.54	1.00
							A/D	6.35	<i>0.04</i>
							B/C	1.47	1.00
							B/D	6.42	<i>0.00</i>
							C/D	7.89	0.06
						χ^2 ; df; <i>p</i> -value	Group	Difference	<i>p</i> -value
FIQ total, median	61.97	67.69	60.72	65.61	51.85	18.13; 3; 0.00	A/B	6.97	<i>0.01</i>
Quartiles	50.39–74.75	55.93–79.78	49.90–72.65	45.54–78.31	35.97–64.99		A/C	2.08	0.57
Min–Max	2.88–97.40	12.70–97.40	19.70–95.47	14.76–95.00	2.88–80.89		A/D	15.84	<i>0.00</i>
							B/C	4.89	0.69
							B/D	8.87	<i>0.01</i>
							C/D	13.76	0.10
FIQ pain, median	7.50	8.00	7.35	7.20	6.60	9.90; 3; 0.019	A/B	0.65	<i>0.05</i>
Quartiles	5.80–8.60	6.30–8.75	5.65–8.50	5.75–8.83	4.90–7.85		A/C	0.80	0.81
Min–Max	0.00–10.00	0.90–10.00	0.20–10.00	3.50–9.20	0.00–9.80		A/D	1.40	<i>0.00</i>
							B/C	0.15	0.85
							B/D	0.75	0.08
							C/D	0.60	0.35
FIQ fatigue, median	8.60	9.00	8.40	8.65	7.90	7.96; 3; 0.047	A/B	0.60	<i>0.03</i>
Quartiles	7.20–9.50	8.10–9.58	7.18–9.33	6.43–9.83	5.43–9.23		A/C	0.35	0.63
Min–Max	1.30–10.00	2.80–10.00	3.00–10.00	1.30–9.90	1.40–10.00		A/D	1.10	<i>0.01</i>
							B/C	0.25	0.87
							B/D	0.50	0.22
							C/D	0.75	0.51
SF-36 PCS, median	26.71	25.78	26.62	30.68	28.63	10.27; 3; 0.016	A/B	0.84	0.42
Quartiles	22.57–30.94	21.87–30.47	22.60–30.51	27.42–34.02	23.03–34.64		A/C	4.90	<i>0.01</i>
Min–Max	8.79–50.68	8.79–40.79	11.02–43.78	20.49–34.02	14.70–50.68		A/D	2.85	<i>0.05</i>
							B/C	4.06	<i>0.01</i>
							B/D	2.01	0.09
							C/D	2.05	0.50
SF-36 MCS, median	40.78	37.43	41.11	43.53	49.04	10.21; 3; 0.017	A/B	3.68	0.43
Quartiles	31.20–50.40	30.31–48.40	30.77–49.73	24.99–50.84	36.70–56.23		A/C	6.10	0.87
Min–Max	14.90–66.59	16.10–64.91	17.26–64.71	14.90–56.73	20.42–66.59		A/D	11.61	<i>0.00</i>
							B/C	2.42	0.73
							B/D	7.93	<i>0.01</i>
							C/D	5.51	0.06

*Group differences in age were analysed using analysis of variance (ANOVA), while groups differences on FIQ and SF-36 were analysed using the Kruskal–Wallis test. Statistically significant group differences were defined as $p < 0.1$.

**Multiple group comparisons in age were performed using Independent samples *t*-test, while multiple group comparisons on FIQ and SF-36 were performed using the Mann–Whitney *U* test.

Statistically significant differences (in italic) in the *post-hoc* analyses were defined at $p < 0.05$.

AMPS: Assessment of Motor and Process Skills; FIQ: Fibromyalgia Impact Questionnaire; FIQ total: overall impact of the fibromyalgia. SF-36: 36-item Short Form; PCS: Physical Composite Score; MCS: Mental Composite Score.

Table II. Group differences in ability to perform activities of daily living (ADL) when evaluated with the Assessment of Motor and Process Skills (AMPS), the 36-item Short Form physical function subscale (SF-36 PF) and the Fibromyalgia Impact Questionnaire physical function subscale (FIQ PF) (n = 257)

	Total	Group A	Group B	Group C	Group D	Group difference* χ^2 ; df; <i>p</i> -value	Significant differences between groups**		
							Group	Difference	<i>p</i> -value
AMPS							A/B	0.15	<i>0.01</i>
Motor, median	1.07	0.91	1.06	1.65	1.66	121.65; 3; 0.00	A/C	0.74	<i>0.00</i>
Quartiles	0.74–1.41	0.48–1.14	0.73–1.27	1.57–1.78	1.56–1.94		A/D	0.75	<i>0.00</i>
Min–Max	0.04–2.82	0.04–1.48	0.06–1.49	1.52–2.02	1.50–2.82		B/C	0.59	<i>0.00</i>
							B/D	0.60	<i>0.00</i>
							C/D	0.01	<i>0.77</i>
AMPS process, median	1.09	0.81	1.23	0.84	1.37	189.74; 3; 0.00	A/B	0.42	<i>0.00</i>
Quartiles	0.85–1.26	0.65–0.89	1.11–1.44	0.77–0.96	1.22–1.58		A/C	0.03	0.13
Min–Max	0.12–2.18	0.12–0.99	1.00–2.18	0.53–0.98	1.05–1.90		A/D	0.56	<i>0.00</i>
							B/C	0.39	<i>0.00</i>
							B/D	0.14	<i>0.01</i>
							C/D	0.53	<i>0.00</i>
FIQ PF, median	5.67	6.25	5.56	5.00	4.51	13.57; 3; 0.00	A/B	0.69	<i>0.02</i>
Quartiles	3.73–7.00	4.46–7.41	3.58–6.67	3.48–7.17	2.00–6.05		A/C	1.25	0.29
Min–Max	0.00–9.00	0.00–9.00	0.00–8.89	0.67–9.00	0.00–8.00		A/D	1.74	<i>0.00</i>
							B/C	0.56	0.92
							B/D	1.05	<i>0.03</i>
							C/D	0.49	<i>0.27</i>
SF-36 PF, median	40.00	40.00	40.00	55.00	55.00	20.82; 3; 0.00	A/B	0.00	0.66
Quartiles	25.00–60.00	25.00–52.50	25.00–55.00	37.50–71.25	40.00–66.25		A/C	15.00	0.24
Min–Max	0.00–95.00	0.00–80.00	0.00–80.00	22.22–85.00	10.00–95.00		A/D	15.00	<i>0.05</i>
							B/C	15.00	0.14
							B/D	15.00	<i>0.01</i>
							C/D	0.00	0.75

*Group differences in AMPS, FIQ PF and SF-36 PF were analysed using the Kruskal–Wallis test and statistically significant group differences were defined as $p < 0.1$.

**Multiple group comparisons in AMPS, FIQ and SF-36 were performed using the Mann–Whitney *U* test.

The statistically significant differences (in italic) in the *post-hoc* analyses were determined by $p < 0.05$ and clinically relevant differences (in bold) were > 0.3 logits in AMPS and > 0.5 SD for the FIQ and SF-36.