



IMPACT OF UNILATERAL SPATIAL NEGLECT WITH OR WITHOUT OTHER COGNITIVE IMPAIRMENTS ON INDEPENDENT GAIT RECOVERY IN STROKE SURVIVORS

Yosuke KIMURA, PT, MSc^{1,2}, Minoru YAMADA, PT, PhD², Daisuke ISHIYAMA, PT, MSc², Naohito NISHIO, PT, MSc², Yota KUNIEDA, PT, MSc², Shingo KOYAMA, PT, MSc², Atushi SATO, PT, MSc², Yuhei OTOBE, PT, MSc², Shunsuke OHJI, PT, MSc², Mizue SUZUKI, ST², Hideyuki OGAWA, PT², Daisuke ITO, OT², Takeo ICHIKAWA, PT², Koji HAMANAKA, PT, MSc¹, Naoki TANAKA, PT¹ and Yasushi MUROH, MD³

From the ¹Department of Rehabilitation and ³Department of Rehabilitation Medicine, Tokyo Shinjuku Medical Center, Japan Community Health care Organization, and ²Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tokyo, Japan

Objective: To investigate the impact of unilateral spatial neglect with or without other cognitive impairments on recovery of independent gait in stroke survivors.

Design: A prospective cohort study.

Subjects: Ninety-four stroke survivors in an inpatient rehabilitation ward.

Methods: The presence of unilateral spatial neglect was assessed by the visuospatial perception score of the Stroke Impairment Assessment Set, and other cognitive impairments were assessed by Mini-Mental State Examination. Participants were categorized into 3 groups: group 1, unilateral spatial neglect with other cognitive impairments; group 2, unilateral spatial neglect without other cognitive impairments; and group 3, non-unilateral spatial neglect. The outcome was the walking score of the Functional Independence Measure (FIM) at discharge (score ≥ 6 or ≤ 5).

Results: Multivariate logistic regression analysis (reference, group 3) showed that the presence of unilateral spatial neglect with other cognitive impairments (group 1) had a significant association with dependence of gait ($p=0.003$), and the odds ratio (95% confidence interval) was 5.55 (1.19–23.04). In contrast, there was no significant relationship between the presence of unilateral spatial neglect without other cognitive impairments (group 2) and dependence of gait ($p=0.207$).

Conclusion: The presence of unilateral spatial neglect without other cognitive impairments is not a significant factor for regaining independent gait. In contrast, unilateral spatial neglect becomes a strong negative factor when combined with other cognitive impairments.

Key words: stroke; hemispatial neglect; cognitive impairment; gait.

Accepted Oct 8, 2018; Epub ahead of print Nov 8, 2018

J Rehabil Med 2019; 51: 00–00

Correspondence address: Yosuke Kimura, Department of Rehabilitation, Tokyo Shinjuku Medical Center, Japan Community Health care Organization, 5-1 Tukudocho, Shinjuku-ku, Tokyo 162-8543, Japan. E-mail: mcbb10ver@yahoo.co.jp

LAY ABSTRACT

Unilateral spatial neglect is characterized by reduced awareness of stimuli on one side of the body. Spatial neglect is a common cognitive impairment in stroke patients with damage to the right side of their brain. In addition, stroke often causes other cognitive impairments, such as memory deficits and non-spatial attention deficits. These neurological symptoms may affect recovery from stroke. This study investigated whether the presence of spatial neglect in combination with other cognitive impairments negatively affects the recovery of gait ability in stroke patients. It was found that, in the absence of other cognitive impairments, spatial neglect did not have a negative impact on recovery of gait ability. In contrast, if associated with other cognitive impairments, spatial neglect was found to exacerbate recovery. This information will be helpful in developing optimal rehabilitation programmes for the recovery of gait ability in stroke patients.

Recovery of independent gait is one of the main goals in rehabilitation of stroke survivors (1). Independence of gait in stroke survivors is associated with ability to perform activities of daily living (ADL) (2), quality of life (3), caregiver burden (4), discharge destination (5) and even prognosis of survival (6). Therefore, it is important to investigate the factors associated with recovery of independent gait in order to develop optimal rehabilitation programmes and discharge plans.

Unilateral spatial neglect (USN) is thought to play a crucial role in the functional disadvantages in stroke survivors. USN leads to significantly poorer recovery of ADL ability (7, 8) and gait dependency (9, 10). However, other previous studies have indicated that the improvement in ADL ability does not differ according to the presence or absence of USN (11), and that the recovery of independent gait is not affected by the severity of USN after controlling for the severity of paresis (12). In addition, relatively recent studies have reported that there are no significant relationships between USN and ability in independent gait in outdoor settings, in chronic stroke survivors (13, 14).

Thus, the association between USN and recovery of gait independence remains controversial.

Other cognitive impairments (CIs), such as memory deficits and non-spatial attention deficits, are common symptoms in stroke survivors, and they negatively influence their functional outcome (15, 16). In addition, previous studies have reported that, compared with stroke survivors without USN, those with USN are more likely to have other cognitive dysfunctions (17, 18). Thus, it is important to consider other cognitive functions in addition to USN in the prediction of gait ability in stroke survivors.

The Mini-Mental State Examination (MMSE) is an easily applicable and most widely used instrument in screening for CIs in stroke survivors. Several studies have reported acceptable validity of the MMSE as a screening tool and its relationship with functional recovery in stroke survivors (19–21). In addition, another previous study reported that the MMSE score reflects the number of disturbed cognitive domains, such as memory, mental speed, and non-spatial attention, in subacute stroke survivors (22). Although the value of the MMSE in screening for cognitive dysfunctions in stroke survivors remains controversial (23), it is considered a useful indicator for conveniently evaluating other cognitive dysfunctions.

This study aimed to investigate the impact of USN with or without other CIs on the recovery of independent gait in subacute stroke survivors. We hypothesized that the presence of USN without other CIs would be less negatively associated with the recovery of independent gait, and that it might be a strong negative factor when combined with other CIs. Clarification of these relationships would help in considering the prognostic prediction and interventions for regaining independent gait.

METHODS

Participants

This prospective cohort study was conducted on stroke survivors admitted to the 37-bed convalescent inpatient rehabilitation ward of our hospital from April 2011 to March 2017. Diagnosis of stroke was based on clinical examination by a physiatrist and an imaging test (computed tomography or magnetic resonance imaging) by a radiologist. Inclusion criteria were: first stroke in the right brain hemisphere; diagnosis of cerebral haemorrhage or cerebral infarction; independence in performing ADL prior to stroke; and requirement for a wheelchair for locomotion on admission. Exclusion criteria were: presence of neuromuscular diseases or severe musculoskeletal diseases, worsening medical conditions during hospitalization (such as a recurrence of stroke or severe infection that would contraindicate rehabilitation), and inability to complete the assessment. In addition, participants who had a diagnosis of dementia prior to stroke, and those who had a pharmacological intervention on cognitive impairments prior to stroke were excluded. The study was conducted in

accordance with the principles of the Declaration of Helsinki, and it was reviewed and approved by the ethics committees of our hospital (approval number: 27–20).

Evaluation of unilateral spatial neglect

The presence of USN was assessed by the visuospatial perception score of the Stroke Impairment Assessment Set (SIAS) (24). The visuospatial perception score of the SIAS was evaluated on admission and at discharge. A 50-cm long tape measure was used for evaluation, and the central point method was adopted. Participants were asked to touch the midpoint of a tape held horizontally 50 cm in front of them, using the unaffected thumb and index finger. Two trials were allowed, and the worst error was used for the scoring value. If there was more than a 15-cm deviation from the midpoint, the score was 0. An error between 15 cm and 5 cm was scored as 1, while an error between 5 cm and 2 cm was scored as 2. A score of 3 meant deviation from the midpoint by less than 2 cm. We defined the presence of USN as a visuospatial perception score of 2 or less. This method was confirmed to have good inter-rater reliability and concurrent validity, assessed via 20-cm line bisection and flower-and-cube copying tasks in stroke survivors (24).

Other cognitive functions

Other cognitive functions were assessed on admission by using the MMSE, which consists of the following 5 areas of cognitive functions: orientation, memory, attention and calculation, language, and construction. The total scores vary from 0 to 30, with higher scores indicating better cognitive functions. In this study, other CIs were defined as a score of less than 24 in the MMSE (the cut-off value was defined by referring to previous studies) (10, 21).

Outcome variable

We investigated the gait dependency of the participants upon discharge, with the walking score of the *Functional Independence Measure (FIM)* (25) as the outcome measure. A FIM walking score of 7 corresponds to complete independence, wherein the participant can safely walk a minimum of 150 ft (50 m) without using assistive devices. A FIM walking score of 6 corresponds to modified independence, wherein the participant can walk a minimum of 150 ft (50 m) without supervision, but with the support of a brace (orthosis) or cane. FIM walking scores of 1–5 correspond to requiring help or supervision and are determined by the level of physical assistance required for walking. In this study, gait independence was defined as a FIM walking score of 6 or more, according to a previous study (26).

Other variables

Demographic characteristics and stroke-related information including age, sex, stroke type (cerebral infarction or cerebral haemorrhage), number of days from onset of stroke to admission, length of stay, body mass index, comorbidity, use of medication (antidepressants and anxiolytics), severity of hemiplegia of the lower limb, and ability to perform ADL were investigated on admission to our rehabilitation ward. Comorbidity was assessed using the Charlson comorbidity index (CCI) (27). The CCI is an evaluation index with 1 to 6 points for 19 comorbidities, with a higher score indicating greater comorbidity. The severity of hemiplegia was determined in terms of the Brunnstrom recovery stages (BRS) (28). The BRS classifies voluntariness in paralysed limbs into 6 ordinal stages, with the lower stages

indicating severe paralysis. ADL ability was assessed using the FIM. The FIM is composed of 18 items divided into 6 subcategories: self-care (6 items), sphincter control (2 items), transfers (3 items), locomotion (2 items), communication (2 items), and social cognition (3 items). Each item is scored on a 7-point ordinal scale from a score of 1 (total dependence) to 7 (complete independence). The high reliability of the FIM for stroke survivors has been reported previously (25).

Rehabilitation treatment

In the Japanese medical insurance system, participants are referred from acute hospitals approximately 30 days after onset of stroke and receive hospital care in convalescent rehabilitation wards for up to 180 days (29). In this study, all participants underwent rehabilitation programmes every day during hospitalization. The programmes were based on a comprehensive approach and included physical, occupational, and speech therapies, as necessary. Participants were provided with approximately 2 h (median 118 (interquartile range (IQR) 100–136) min) of rehabilitation programmes per day. There was no specific protocol or procedure for treating USN in this study. All participants with USN underwent approximately equivalent amounts of conventional therapies, such as visual scanning training, trunk rotation training, and feedback training in ADL tasks. In addition, participants with other CIs underwent conventional cognitive training.

Data analysis

The participants were assigned to 3 groups according to the presence or absence of USN and the MMSE score on admission:

group 1, USN with other CIs; group 2, USN without other CIs; and group 3, non-USN (Fig. 1).

The characteristics of the participants were compared across the groups by 1-way analysis of variance, Kruskal–Wallis test, χ^2 test, and Fisher's exact test after evaluating the normality of the variables using the Shapiro–Wilk test. To examine the effect of the USN and other CIs on recovery of independent gait, we used logistic regression analysis using the 3 groups as the independent variables (reference, group 3) and the FIM walking score (score ≥ 6 or ≤ 5) as the dependent variables. In the logistic regression analysis, 2 models were used. In the first model, we did not adjust for covariates (Model 1, Crude); in the second model, in addition to Model 1, variables with $p < 0.05$ in univariate analysis were included as covariates (Model 2, multivariate model).

Statistical significance was defined as a p -value less than 0.05 for all analyses. Statistical analyses were performed using the SPSS software version 24.0 (IBM, Tokyo, Japan).

RESULTS

During the study period, 131 consecutive stroke survivors met the inclusion criteria, and 94 were analysed in the present study (Fig. 1).

The characteristics of the participants are shown in Table I. The mean age of the study participants was 69.9 years \pm 9.3, and 57 (62.8%) were men. A total of 44 participants (46.8%) had had an ischaemic stroke




Fig. 1. Flow chart of the participants' selection process. USN: unilateral spatial neglect; CIs: cognitive impairments; MMSE: Mini-Mental State Examination.

and 50 (53.2%) had had a haemorrhagic stroke. Group 1 (USN with other CIs) consisted of 30 participants (31.9%); group 2 (USN without other CI), 26 participants (27.7%); and group 3 (non-USN), 38 participants (40.4%). With regard to baseline characteristics, significant differences in the length of stay ($p=0.002$), BRS of the lower limb ($p=0.033$), and FIM scores on admission ($p<0.001$) were observed among the 3 groups. The prevalence of USN at discharge was 76.7% ($n=23$) in group 1 and 53.8% ($n=14$) in group 2 ($p=0.072$). The frequency of regaining independent gait upon discharge was 35.1% ($n=33$) overall, 10.0% ($n=3$) in group 1, 50.0% ($n=13$) in group 2, and 44.7% ($n=17$) in group 3 ($p<0.001$).

The results of the logistic regression analysis for the 3 groups (reference, group 3) in regaining independent gait are shown in Table II. In Model 1 (Crude model), the presence of USN with other CIs (group 1) showed a significant association with dependence of gait upon discharge ($p=0.004$), and the odds ratio (OR) (95% confidence interval (95% CI)) for dependence of gait was 7.29 (1.18–28.20). In contrast, there was a non-significant relationship between the presence of USN without other CIs (group 2) and dependence of gait at discharge ($p=0.679$). After adjustment for covariates (Model 2), the presence of USN with other CIs (group 1) still showed a significant association with dependence of gait ($p=0.003$), and the OR (95% CI) for dependence of gait was 5.55 (1.19–23.04). There was a non-significant relationship between the presence of USN without other CIs (group 2) and dependence of gait at discharge ($p=0.207$).

Table II. Logistic regression analysis for dependence of gait at discharge.

Factors	Model 1		Model 2	
	OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>
Groups according to USN and other cognitive impairments				
Group 1	7.29 (1.18–28.20)	0.004	5.55 (1.19–23.04)	0.003
Group 2	0.81 (0.29–2.20)	0.679	2.06 (0.58–8.27)	0.207
Group 3	1.00 ref.		1.00 ref.	
Age			1.07 (0.99–1.14)	0.065
Motor FIM on admission			0.91 (0.85–0.97)	0.002
Brunnstrom recovery stage of lower extremities				
Severe (1–2)			1.44 (0.25–8.36)	0.687
Moderate (3–4)			0.43 (0.11–1.71)	0.232
Mild (5–6)			1.00 ref.	

Group 1: USN with other CIs; Group 2: USN without other CIs; Group 3: non-USN. Model 1: Crude model. Model 2: Multivariate model adjusted for covariates that $p<0.05$ in univariate analysis. OR: odds ratio; 95% CI: 95% confidence interval; ref: reference; USN: unilateral spatial neglect; FIM: Functional Independence Measure.

DISCUSSION

Right hemisphere stroke survivors were assigned to 3 groups based on the presence or absence of USN and on MMSE score at admission to the convalescent inpatient rehabilitation ward and their association with recovery of independent gait was investigated. The results showed that the presence of USN without other CIs (group 2) was not significantly related to recovery of independent gait during hospitalization. However, the presence of USN with other CIs (group 1) had a significant negative impact on the recovery of independent gait, even after adjustment for covariates such as age, motor-FIM score on admission, and severity of hemiplegia of the lower limb. Thus, the presence of USN becomes a strong negative predictor of independent gait recovery when combined with other cognitive dysfunctions.

Table I. Characteristics of the participants

	Overall (<i>n</i> = 94)	Group 1 (<i>n</i> = 30)	Group 2 (<i>n</i> = 26)	Group 3 (<i>n</i> = 38)	<i>p</i>
Age, years, mean (SD)	69.9 (9.3)	71.1 (10.4)	66.8 (8.6)	71.1 (8.6)	0.135
Gender, men, <i>n</i> (%)	57 (62.8)	17 (56.7)	20 (76.9)	22 (57.9)	0.357
BMI, kg/m ² , mean (SD)	22.2 (3.7)	21.7 (3.8)	22.0 (3.0)	22.5 (3.7)	0.667
Type of stroke (cerebral infraction), <i>n</i> (%)	44 (46.8)	14 (46.7)	8 (30.8)	22 (57.9)	0.102
Days between stroke onset and admission, median (IQR)	32.0 (22.0–43.5)	36.5 (28.3–45.5)	33.0 (24.8–48.0)	30.0 (17.8–42.0)	0.121
Length of stay, days, median (IQR)	140.5 (103.3–165.0)	162.5 (123.3–170.0)	160.0 (130.0–170.3)	134.5 (105.0–148.0)	0.002
Charlson comorbidity index, median (IQR)	2.0 (2.0–3.0)	2.0 (2.0–3.0)	2.0 (2.0–3.0)	3.0 (2.0–4.0)	0.105
Medication					
Antidepressant, <i>n</i> (%)	9 (9.6)	3 (10.0)	5 (19.2)	1 (2.6)	0.082
Anxiolytic, <i>n</i> (%)	22 (23.4)	8 (26.7)	9 (34.6)	5 (13.2)	0.113
Brunnstrom recovery stage of lower extremities					
Severe (1–2), <i>n</i> (%)	28 (29.8)	15 (30.0)	7 (26.9)	6 (15.8)	
Moderate (3–4), <i>n</i> (%)	43 (45.7)	11 (36.7)	13 (50.0)	19 (50.0)	
Mild (5–6), <i>n</i> (%)	23 (24.5)	4 (13.3)	6 (23.1)	13 (34.2)	
MMSE score on admission, points, median (IQR)	25.0 (19.3–28.0)	19.0 (15.0–21.8)	28.0 (26.0–28.0)	28.0 (21.0–29.0)	<0.001
FIM on admission, points, mean (SD)	59.6 (14.9)	50.1 (14.1)	65.1 (12.9)	63.3 (13.4)	<0.001
Motor FIM on admission, points, mean (SD)	36.9 (11.6)	31.0 (11.7)	40.1 (11.3)	39.4 (10.2)	0.002
Cognitive FIM on admission, points, mean (SD)	22.6 (4.9)	19.0 (5.2)	25.0 (2.5)	23.8 (4.4)	<0.001
Walk item of FIM on admission, points, median (IQR)	1.0 (1.0–1.0)	1.0 (1.0–1.0)	1.0 (1.0–1.0)	1.0 (1.0–1.0)	0.687
Walk item of FIM at discharge, points, median (IQR)	5.0 (1.0–6.0)	1.0 (1.0–3.5)	5.0 (5.0–6.0)	5.0 (3.0–6.0)	<0.001
Presence of USN at discharge, yes, <i>n</i> (%)	37 (39.4)	23 (76.7)	14 (53.8)	–	0.072
Gait ability at discharge, independence, yes, <i>n</i> (%)	33 (35.1)	3 (10.0)	13 (50.0)	17 (44.7)	0.002

Group 1: USN with other CIs; Group 2: USN without other CIs; Group 3: non-USN. Each variables were compared among three groups by Analysis of variance, Kruskal-Wallis test, Chi-square test, and Fisher's exact test, respectively. Only the presence of USN at discharge was compared between group 1 and group 2. SD: standard deviation; IQR: interquartile range; BMI: body mass index; MMSE: Mini-Mental State Examination; FIM: Functional Independence Measure; USN: unilateral spatial neglect; CIs: cognitive impairments.

These results showed that USN was not significantly associated with recovery of independent gait in stroke survivors without other CIs. This result is consistent with some previous studies, which showed that USN had no negative prognostic influence on gait ability and ADL ability (12, 30). Several previous studies have investigated the time course of USN recovery and shown that the severity of neglect greatly improved until about 12 weeks after the onset of stroke (31–33). Another recent study reported that the improvement in the severity of neglect significantly correlated with increased MMSE scores in subacute stroke survivors (34). In our results also, neglect symptoms disappeared at discharge in approximately half of the stroke survivors in group 2. Therefore, the influence of the presence of USN at admission on independent gait recovery might be limited in stroke survivors with relatively good other cognitive functions. In contrast, approximately 75% of stroke survivors in group 1 still had neglect symptoms at discharge and were significantly inhibited from regaining independent gait.

Several studies have concluded that USN is a negative predictive factor of functional outcome in stroke survivors (7–10, 34, 35). Unlike these previous studies, our results showed that USN was not significantly associated with recovery of independent gait in stroke survivors without other CIs. Stroke survivors with USN often have other cognitive dysfunctions that negatively influence functional recovery. However, some of the previous studies that concluded USN as a negative predictive factor of functional outcome did not consider other cognitive functions (7, 34, 35), in other words, whether USN itself or its combination with other CIs played a crucial role in functional recovery is unclear. The influence of USN may be overestimated unless other cognitive functions are considered.

The main finding of this study is that the presence of USN with other CIs had a strong negative impact on recovery of independent gait. Stroke survivors with USN are unable to orient their attention toward the left hemi-space and often are not aware of the left side of their body while performing everyday tasks. These neglect symptoms could lead to unstable walking, for instance, because stroke survivors with spatial neglect tend to bump into objects (36). Awareness regarding disability is a key determinant to overcoming these neglect symptoms in stroke survivors with USN (37). Cognitive skills, such as short-term verbal memory, non-spatial attention, comprehension, and orientation, are fundamental to the awareness and understanding of the impairments, as well as understanding the relationship between insight and the capacity to learn. Deficits in these cognitive skills could interfere with stroke survivors' awareness of neglect symptoms

and inhibit learning of compensatory strategies to overcome disabilities in the rehabilitation and ADL settings. Therefore, the presence of USN with a low MMSE score may have a strongly negative impact on recovery of independent gait in stroke survivors.

Study limitations

This study has several limitations. First, the subtypes of USN were not considered. A previous study reported that the improvement in ADL differed depending on the subtypes of USN (38), and that it could also be associated with gait independence. Therefore, it is necessary to consider a more nuanced definition of USN in future studies. Secondly, the sample size was relatively small; in particular, only 3 participants in group 1 regained independent gait. Further studies will require a larger cohort and the inclusion of stroke survivors with various severities. Thirdly, we did not examine the more detailed domains of cognitive functions, such as memory, language, attention, and executive function. Further studies are warranted to investigate the relationships between specific domains of cognitive impairment and functional outcomes. Finally, we did not investigate information regarding apathy and anosognosia for hemiplegia. Apathy and anosognosia are often found in stroke survivors and are known to have a negative effect on functional outcome (30, 39, 40). Further studies should investigate these symptoms and take into account their influence on the recovery of gait ability in stroke survivors.

Conclusion

The presence of USN in subacute stroke survivors with relatively good other cognitive functions was not significantly associated with recovery of independent gait during hospitalization in the convalescent rehabilitation ward. In contrast, stroke survivors with USN and other CIs were significantly inhibited from regaining independent gait even after controlling for covariates such as ADL ability at admission, and severity of hemiplegia. These findings indicated that USN became a strong negative predictor when combined with other cognitive dysfunctions, and that only 10% of the stroke survivors with both USN and other CIs regained independent gait during hospitalization. These results may be helpful in accurate prediction of the prognosis of subacute stroke and decisions regarding interventions required for recovery of independent gait in subacute stroke survivors.

ACKNOWLEDGEMENTS

We thank the nursing staff and rehabilitation therapists of JCHO Tokyo Shinjuku Medical Center for their contributions to the data collection.

We are also grateful to our laboratory members and Ms Mai Kawamura for their helpful advice on drafts of this manuscript.

REFERENCES

- Dobkin BH. Clinical practice. Rehabilitation after stroke. *N Engl J Med* 2005; 352: 1677–1684.
- Cho KH, Lee JY, Lee KJ, Kang EK. Factors related to gait function in post-stroke patients. *J Phys Ther Sci* 2014; 26: 1941–1944.
- Kyung Kim, Young Mi Kim, Eun Kyung Kim. Correlation between the activities of daily living of stroke patients in a community setting and their quality of life. *J Phys Ther Sci* 2014; 26: 417–419.
- Vincent C, Desrosiers J, Landreville P, Demers L; BRAD group. Burden of caregivers of people with stroke: evolution and predictors. *Cerebrovasc Dis* 2009; 27: 456–464.
- Brauer SG, Bew PG, Kuys SS, Lynch MR, Morrison G. Prediction of discharge destination after stroke using the motor assessment scale on admission: a prospective, multisite study. *Arch Phys Med Rehabil* 2008; 89: 1061–1065.
- Chiu HT, Wang YH, Jeng JS, Chen BB, Pan SL. Effect of functional status on survival in patients with stroke: is independent ambulation a key determinant? *Arch Phys Med Rehabil* 2012; 93: 527–531.
- Di Monaco M, Schintu S, Dotta M, Barba S, Tappero R, Gindri P. Severity of unilateral spatial neglect is an independent predictor of functional outcome after acute inpatient rehabilitation in individuals with right hemispheric stroke. *Arch Phys Med Rehabil* 2011; 92: 1250–1256.
- Paolucci S, Antonucci G, Grasso MG, Pizzamiglio L. The role of unilateral spatial neglect in rehabilitation of right brain-damaged ischemic stroke patients: a matched comparison. *Arch Phys Med Rehabil* 2001; 82: 743–749.
- Paolucci S, Bragoni M, Coiro P, De Angelis D, Fusco FR, Morelli D, et al. Quantification of the probability of reaching mobility independence at discharge from a rehabilitation hospital in nonwalking early ischemic stroke patients: a multivariate study. *Cerebrovasc Dis* 2008; 26: 16–22.
- Nijboer T, van de Port I, Schepers V, Post M, Visser-Meily A. Predicting functional outcome after stroke: the influence of neglect on basic activities in daily living. *Front Hum Neurosci* 2013; 7: doi: 10.3389/fnhum.2013.00182.
- Chen P, Hreha K, Kong Y, Barrett AM. Impact of spatial neglect on stroke rehabilitation: evidence from the setting of an inpatient rehabilitation facility. *Arch Phys Med Rehabil* 2015; 96: 1458–1466.
- van Nes IJ, van Kessel ME, Schils F, Fasotti L, Geurts AC, Kwakkel G. Is visuospatial hemineglect longitudinally associated with postural imbalance in the postacute phase of stroke? *Neurorehabil Neural Repair* 2009; 23: 819–824.
- Durcan S, Flavin E, Horgan F. Factors associated with community ambulation in chronic stroke. *Disabil Rehabil* 2016; 38: 245–249.
- Ferreira MS, Chamlian TR, França CN, Massaro AR. Non-motor factors associated with the attainment of community ambulation after stroke. *Clin Med Res* 2015; 13: 58–64.
- Park J, Lee SU, Jung SH. Prediction of post-stroke functional mobility from the initial assessment of cognitive function. *NeuroRehabilitation* 2017; 41: 169–177.
- Robertson IH, Ridgeway V, Greenfield E, Parr A. Motor recovery after stroke depends on intact sustained attention: a 2-year follow-up study. *Neuropsychology* 1997; 11: 290–295.
- Lee BH, Kim EJ, Ku BD, Choi KM, Seo SW, Kim GM, et al. Cognitive impairments in patients with hemispatial neglect from acute right hemisphere stroke. *Cogn Behav Neurol* 2008; 21: 73–76.
- Linden T, Samuelsson H, Skoog I, Blomstrand C. Visual neglect and cognitive impairment in elderly patients late after stroke. *Acta Neurol Scand* 2005; 111: 163–168.
- Tombaugh TN, McIntyre NJ. The Mini-Mental State Examination: a comprehensive review. *J Am Geriatr Soc* 1992; 40: 922–935.
- Zwecker M, Levenkrohn S, Fleisig Y, Zeilig G, Ohry A, Adunsky A. Mini-Mental State Examination, cognitive FIM instrument, and the Loewenstein Occupational Therapy Cognitive Assessment: relation to functional outcome of stroke patients. *Arch Phys Med Rehabil* 2002; 83: 342–345.
- Patel MD, Coshall C, Rudd AG, Wolfe CD. Cognitive impairment after stroke: clinical determinants and its associations with long-term stroke outcomes. *J Am Geriatr Soc* 2002; 50: 700–706.
- Bour A, Rasquin S, Boreas A, Limburg M, Verhey F. How predictive is the MMSE for cognitive performance after stroke? *J Neurol* 2010; 257: 630–637.
- Burton L, Tyson SF. Screening for cognitive impairment after stroke: a systematic review of psychometric properties and clinical utility. *J Rehabil Med* 2015; 47: 193–203.
- Liu M, Chino N, Tuji T, Masakado Y, Hase K, Kimura A. Psychometric properties of the Stroke Impairment Assessment Set (SIAS). *Neurorehabil Neural Repair* 2002; 16: 339–351.
- Ottenbacher KJ, Hsu Y, Granger CV, Fiedler RC. The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil* 1996; 77: 1226–1232.
- Patel AT, Duncan PW, Lai SM, Studenski S. The relation between impairments and functional outcomes poststroke. *Arch Phys Med Rehabil* 2000; 81: 1357–1363.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987; 40: 373–383.
- Brunnstrom S. Motor testing procedures in hemiplegia: based on sequential recovery stages. *Phys Ther* 1966; 46: 357–375.
- Miyai I, Sonoda S, Nagai S, Takayama Y, Inoue Y, Kakehi A, et al. Results of new policies for inpatient rehabilitation coverage in Japan. *Neurorehabil Neural Repair* 2011; 25: 540–447.
- Pedersen PM, Jørgensen HS, Nakayama H, Raaschou HO, Olsen TS. Hemineglect in acute stroke – incidence and prognostic implications. The Copenhagen Stroke Study. *Am J Phys Med Rehabil* 1997; 76: 122–127.
- Ringman JM, Saver JL, Woolson RF, Clarke WR, Adams HP. Frequency, risk factors, anatomy, and course of unilateral neglect in an acute stroke cohort. *Neurology* 2004; 63: 468–474.
- Nijboer TC, Kollen BJ, Kwakkel G. Time course of visuospatial neglect early after stroke: a longitudinal cohort study. *Cortex* 2013; 49: 2021–2027.
- Cassidy TP, Lewis S, Gray CS. Recovery from visuospatial neglect in stroke patients. *J Neurol Neurosurg Psychiatry* 1998; 64: 555–557.
- Cumming TB, Plummer-D'Amato P, Linden T, Bernhardt J. Hemispatial neglect and rehabilitation in acute stroke. *Arch Phys Med Rehabil* 2009; 90: 1931–1936.
- Cherney LR, Halper AS, Kwasnica CM, Harvey RL, Zhang M. Recovery of functional status after right hemisphere stroke: relationship with unilateral neglect. *Arch Phys Med Rehabil* 2001; 82: 322–328.
- Tromp E, Dinkla A, Mulder T. Walking through doorways: an analysis of navigation skills in patients with neglect. *Neuropsychol Rehabil* 1995; 5: 319–331.
- Tham K, Borell L, Gustavsson A. The discovery of disability: a phenomenological study of unilateral neglect. *Am J Occup Ther* 2000; 54: 398–406.
- Spaccavento S, Cellamare F, Falcone R, Loverre A, Nardulli R. Effect of subtypes of neglect on functional outcome in stroke patients. *Ann Phys Rehabil Med* 2017; 60: 376–381.
- Appelros P, Karlsson GM, Seiger A, Nydevik I. Neglect and anosognosia after first-ever stroke: incidence and relationship to disability. *J Rehabil Med* 2002; 34: 215–220.
- Caeiro L, Ferro JM, Costa J. Apathy secondary to stroke: a systematic review and meta-analysis. *Cerebrovasc Dis* 2013; 35: 23–39.